

New study explains near-annual Monsoon oscillations generated by El Niño


October 19, 2015

A new research study by a team of climate researchers from the University of Hawai‘i at Mānoa explains for the first time the source of near-annual pressure and wind changes discovered previously in the Southeast Asian Monsoon system.

The results, published this week in the journal *Proceedings of the National Academy of Sciences*, show how the El Niño phenomenon interacts with the annual cycle of solar radiation in the western Pacific to generate a suite of new atmospheric pressure oscillations that affects wind and rainfall patterns in one Southeast Asia, one of the densest populated areas on our planet.

There still remain major uncertainties about how the atmospheric circulation and the rainfall patterns over Southeast Asia and the Western Tropical Pacific respond to El Niño conditions, such as the current 2015 event. The new findings by Malte Stuecker and Fei-Fei Jin, from the Department of Atmospheric Sciences, and Axel Timmermann from the International Pacific Research Center at the University of Hawai‘i, Mānoa demonstrate that the atmospheric reaction is much more predictable than previously assumed.

September 2015 sea surface temperature anomalies



Current El Niño conditions: Sea surface temperature anomalies [in °Celsius] in September 2015 from NOAA’s Extended Reconstructed Sea Surface Temperature (ERSST) dataset.

As a result of the developing and decaying El Niño conditions and the seasonal march of the sun across the equator, a series of near-annual atmosphere oscillations is generated with periods of about 24, 16, 10, and 8 months, each with its own characteristic pressure pattern. All of these contribute to the formation of an extended and very persistent high pressure system over the western tropical Pacific, which peaks in January and re-emerges in the boreal summer of the following La Niña year.

“Known as the Philippine Sea Anticyclone, this pattern plays a pivotal role in how the effects of El Niño are expressed in Southeast Asia,” explains Stuecker, climate scientist and lead author of the study. “In the past, many researchers looked at seasonal averages of wind and rainfall and missed the details of this variability,” he adds.

“Our new theory also explains the observed persistence of the Philippine Sea Anticyclone, even months after El Niño warming has subsided and well into the subsequent La Niña summer. The effect of El Niño in winter is qualitatively similar to the effect of La Niña in summer. These affects occur when the phases of Pacific warming and the annual solar cycle coincide,” says co-author Fei-Fei Jin, climate researcher and Professor at the Department of Atmospheric Sciences.

“These near-annual monsoon oscillations have some resemblance to the different ringing tones of a bell. Finding a new structure in the chaos of otherwise random weather variability has been very exciting, because it may open the door to enhanced seasonal predictability,” says co-author Axel Timmermann, Professor at the Oceanography Department and the International Pacific Research Center.

The scientists emphasize that this mechanism provides a fundamentally new way to understand variability in the atmosphere on a large range of timescales and can be applied to a number of different climate phenomena.

Citation: Stuecker, M.F., F.-F. Jin, and Timmermann, A.: El Niño-Southern Oscillation frequency cascade. *Proceedings of the National Academy of Sciences*: www.pnas.org/cgi/doi/10.1073/pnas.1508622112 (2015)

Funding Source: M.F.S. and F-F.J. were supported by US National Science Foundation (NSF) Grant AGS-1406601 and US Department of Energy Grant DE-SC0005110. A.T. was supported through NSF grant No. 1049219.

Author Contacts:

Malte F. Stuecker: stuecker@soest.hawaii.edu; +1-808-956-7110

Fei-Fei Jin: jff@hawaii.edu; +1-808-956-4645

Axel Timmermann: axel@hawaii.edu; Tel: +1-808-956-2720

IPRC Media Contact: Rachel Lentz, rlentz@hawaii.edu, +1-808-956-8175

The International Pacific Research Center (IPRC) of the School of Ocean and Earth Science and Technology (SOEST) at the University of Hawai‘i at Mānoa, is a climate research center founded to gain greater understanding of the climate system and the nature and causes of climate variation in the Asia-Pacific region and how global climate changes may affect the region. Established under the "U.S.-Japan Common Agenda for Cooperation in Global Perspective" in October 1997, the IPRC is a collaborative effort between agencies in Japan and the United States.